



Application Procedures

Use Oxivir® Five 16 Concentrate One Step Disinfectant Cleaner to clean and disinfect hard, non-porous surfaces.




Apply use solution. Let stand for 5 minutes. Wipe or allow to air dry.

Use Virex® Plus One-Step Disinfectant Cleaner and Deodorant to clean restroom toilets, sinks, floors and walls.




Apply use-solution to hard, nonporous surfaces. Thoroughly wet surface with a cloth, mop, sponge, sprayer or by immersion. Allow to remain wet for 3 minutes. Wipe dry or allow to air dry.

Apply Avert Sporicidal Disinfectant Cleanersolution by spraying evenly onto entire surface, or by pre-wetting a reusable or disposable microfiber cleaning cloth.



Pre-clean heavily soiled areas. Apply the solution evenly over the entire surface and wait for the appropriate contact time (1 minute for most common microorganisms, 4 minutes for C. difficile). All surfaces must remain visibly wet for 1 minute (4 minutes for C. difficile). If required, wipe surfaces with a clean, damp cloth and let air dry.

Use Oxivir® Tb one-step cleaner disinfectant to wipe hard, non-porous surfaces.



Pull wipe from canister, wipe surfaces, allowing them to stay wet for one minute. Allow to air dry or rinse according to use instructions on some surfaces.

Questions: 1-800-558-2332

Always wear personal protective equipment.